



FERENCE

Section 24(c) Special Local Need Label

**FOR USE IN KINGS, QUEENS, NASSAU AND SUFFOLK COUNTIES IN NEW YORK STATE
ONLY ON GOLF COURSE TURF**

**Ference®
For Control of Annual Bluegrass Weevil on Golf Course Turfgrass Only**

**EPA Reg. No. 100-1551
EPA SLN No. NY-190010**

**This label expires and must not be distributed or used in accordance with
this SLN registration after December 31, 2024**

KEEP OUT OF REACH OF CHILDREN

DIRECTIONS FOR USE

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This label must be in the possession of the user at the time of application.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA-registered label.

The following restriction is required to permit use of Ference in the State of New York:

- Product is restricted use by certified applicators only in New York State

APPLICATION RATES FOR GOLF COURSES

For maximum residual effectiveness or for optimal control of large pest infestations, apply Ference at up to 24 fluid ounces per acre (= 0.55 fluid ounces per 1,000 square feet or 0.313 lb AI/A) to control any of the pests listed below. Apply Ference in sufficient water to provide optimal distribution in the treated area. Use properly calibrated application equipment that will produce a uniform, coarse droplet spray, using a low pressure setting to eliminate off target drift.

Restriction: Do not apply more than 32 fluid ounces (equivalent to 0.42 lb of active ingredient) of product per acre per year in broadcast applications to turfgrass.

APPLICATION TIMING

Annual Bluegrass Weevil: Apply Ference when overwintered adult annual bluegrass weevils are observed and early instar larvae are hatching and penetrating into turfgrass sheaths. The higher rates listed in Table 1 may be required to control late instar larvae.

Ference also provides control of summer annual bluegrass weevil generations. Apply Ference when adults are observed and early instar larvae are hatching and penetrating into turfgrass sheaths. The higher rates listed in Table 1 may be required to control late instar larvae.

Use Ference as part of a program to prevent turf damage from overwintering and following generations of annual bluegrass weevil larvae. Consult your local Syngenta representative, cooperative extension service specialist or pest control advisor for the latest information on using Ference to control annual bluegrass weevil larvae.

Table 1: Turf Application Rates

Target Pest	Ference Turf Application Rates		
	Product (fl oz) per Acre	Product (fl oz) per 1,000 sq ft	Ib A/A
Annual bluegrass weevil	12 to 20	0.275 to 0.459	0.157 to 0.261

©2019 Syngenta

Ference® and the Syngenta logo are trademarks of a Syngenta Group Company

24(c) Registrant:
Syngenta Crop Protection, LLC
P. O. Box 18300
Greensboro, NC 27419-8300

Label Code: NY1551079AA1119